

Spotting imprints of dark matter in the extragalactic Fermi sky with photon counts statistics

Hannes-S. Zechlin



Istituto Nazionale di Fisica Nucleare (INFN), Turin, Italy

Silvia Manconi, Fiorenza Donato

Università di Torino and INFN, Turin, Italy

Gamma-ray searches for dark matter (DM) are often driven by investigating the composition of the extragalactic gamma-ray background (EGB). Statistical methods have recently been proven to outperform the sensitivity of classic approaches in finding unresolved point-source populations and EGB decomposition. We employ the 1-point photon count statistics of eight years of Fermi data to decompose the EGB for latitudes $|b| > 30$ deg, between 1 and 10 GeV. We extend the analysis to incorporate a potential contribution from annihilating DM. Given different interstellar emission models, we set upper bounds on the DM self-annihilation cross section which are strikingly competitive with constraints obtained by other indirect detection methods.


- ▶ statistical analysis (1-point PDF)
- ▶ 8 yrs Pass 8 data, 1 to 10 GeV
- ▶ high Galactic latitudes
- ▶ smooth Galactic DM halo
- ▶ upper limits on DM strikingly competitive with dSphs



Zechlin, Manconi, Donato, arXiv:1710.01506

I. Introduction and 1pPDF [1-4]

We consider the celestial region of interest (ROI) to be partitioned into N_{pix} pixels of equal area Ω_{pix} . The probability p_k of finding k photons in a given pixel is by definition the 1-point probability distribution function (1pPDF). In the simplest scenario of purely isotropic emission, p_k follows a Poisson distribution with an expectation value equal to the mean photon rate. The imprints of more complex diffuse components and a distribution of point sources alter the shape of the 1pPDF, in turn allowing us to investigate these components by measuring the 1pPDF of the data.



- 1pPDF can be modeled with probability generating functions

$$\mathcal{P}^{(p)}(t) = \sum_{k=0}^{\infty} p_k^{(p)} t^k, \quad p_k^{(p)} = \frac{1}{k!} \left. \frac{d^k \mathcal{P}^{(p)}(t)}{dt^k} \right|_{t=0}$$

- **Model for the high-latitude gamma-ray sky**

- isotropic distribution of gamma-ray point sources (dN/dS)
 - > multiply broken power law (MBPL); **parameters freely adjustable**
- diffuse component of Galactic foreground emission
 - > official Fermi template [5]; models A, B, C from [6]; **free normalisation A_{gal}**
- diffuse isotropic background emission
 - > power law (index 2.3); free normalisation
- smooth distribution of Galactic DM
 - > Galactic DM halo, Einasto profile with $\rho(r_{\odot}) = 0.4 \text{ GeV cm}^{-3}$; **free normalisation A_{DM}**

- **pixel-dependent likelihood function**

(full exploitation of spatial templates)

$$\mathcal{L}(\Theta) = \prod_{p=1}^{N_{\text{pix}}} P(k_p), \quad \text{where } P(k_p) \text{ is given by the } p_k^{(p)} \text{ coefficients}$$

In this way, qualitatively, diffuse components are treated as classic template fits, while a distribution of point sources, dN/dS , adds non-Poissonian components.

- **parameter estimation**

—> profile likelihood from Bayesian posterior (MCMC sampling: MultiNest)

- **data set**

—> **Fermi-LAT**: Pass 8, 8 years, 1 to 10 GeV (3 energy bins), UCV, PSF3

—> **ROI**: $|\text{bl}| > 30$ deg, with Fermi Bubbles and Galactic Loop I masked

- **analysis objective**

—> investigate 1pPDF sensitivity reach for additional DM component

—> provide upper limits (ULs) on DM self-annihilation cross section $\langle\sigma v\rangle$, given $A_{\text{DM}} \propto \langle\sigma v\rangle$


- **Galactic foreground (GF) systematics**

—> GF models equipped with high systematic uncertainties

—> possible dependences on or degeneracies of the DM component with GF (in particular with inverse Compton emission) need to be accounted for properly

—> issues mitigated by focusing on high Galactic latitudes only, and ROI optimisation





—> systematic uncertainties of ULs estimated using 4 different GF models



Integrated Galactic foreground emission between 1.99 and 5.0 GeV in the considered ROI.

II. Results [4]

- **upper limits obtained using the official Fermi GF model and models A, B, C**
- moderate systematic scatter
- ➔ **ULs strikingly competitive with bounds recently obtained from dSphs**



References

- [1] Malyshev, D., Hogg, D.W. (2011), ApJ 738, 181
- [2] Zechlin, H.-S., Cuoco, A., Donato, F., et al. (2016), ApJS 225, 18
- [3] Zechlin, H.-S., Cuoco, A., Donato, F., et al. (2016), ApJL 826, L31
- [4] Zechlin, H.-S., Manconi, S., Donato, F. (2017), arXiv:1710.01506
- [5] Acero, F., Ackermann, M., Ajello, M., et al. (2016), ApJS 223, 26
- [6] Ackermann, M., Ajello, M., Albert, A., et al. (2015), ApJ 799, 86